



# Ecological network metrics: opportunities for synthesis

## Citation

Lau, Matthew K., Stuart R. Borrett, Benjamin Baiser, Nicholas J. Gotelli, and Aaron M. Ellison. 2017. "Ecological Network Metrics: Opportunities for Synthesis." *Ecosphere* 8 (8) (August): e01900. doi:10.1002/ecs2.1900.

## Published Version

doi:10.1002/ecs2.1900

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:34389679>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

# Ecological Network Metrics: Opportunities for Synthesis

Matthew K. Lau<sup>1,\*</sup>, Stuart R. Borrett<sup>2,3</sup>, Benjamin Baiser<sup>4</sup>,  
Nicholas J. Gotelli<sup>5</sup>, Aaron M. Ellison<sup>1</sup>

1. Harvard Forest, Harvard University; 2. Department of Biology and Marine Biology, University of North Carolina Wilmington; 3. Duke Network Analysis Center, Social Science Research Institute, Duke University; 4. Department of Wildlife Ecology and Conservation, University of Florida; 5. Department of Biology, University of Vermont; \*corresponding author: Matthew K. Lau (matthewklau@fas.harvard.edu)

## Abstract

Network ecology provides a systems basis for approaching ecological questions, such as factors that influence biological diversity, the role of particular species or particular traits in structuring ecosystems, and long-term ecological dynamics (e.g. stability). Whereas the introduction of network theory has enabled ecologists to quantify not only the degree, but also the architecture of ecological complexity, these advances have come at the cost of introducing new challenges, including new theoretical concepts and metrics, and increased data complexity and computational intensity. Synthesizing recent developments in the network ecology literature, we point to several potential solutions to these issues: integrating network metrics and their terminology across sub-disciplines; benchmarking new network algorithms and models to increase mechanistic understanding; and improving tools for sharing ecological network research, in particular “model” data provenance, to increase the reproducibility of network models and analyses. We propose that applying these solutions will aid in synthesizing ecological subdisciplines and allied fields by improving the accessibility of network methods and models.

*Keywords:* Network ecology, systems analysis, computational methods, metrics, benchmarking, data provenance

# 1 Introduction

Interactions are at the heart of ecology and drive many of its key questions. What are the roles of species interactions in ecological systems? When and why is biological diversity important? What factors influence the long-term dynamics of ecosystems? These are all questions with a long history in ecology (Cherrett, 1989; Council, 2003; Lubchenco et al., 1991; Sutherland et al., 2013) that are not addressed in isolation. Points of intersection include the relationship between diversity and stability (May, 2001, 2006); the identity and role of species that are the main drivers of community structure (Paine, 1966, e.g. keystone species), ecosystem engineers (Jones et al., 1994), or foundation species (Dayton, 1972; Ellison et al., 2005); and the causes and consequences of introducing new species into existing assemblages (Baiser et al., 2008; Simberloff and Holle, 1999). Furthermore, “systems thinking” has been a persistent thread throughout the history of ecology (Margalef, 1963; Odum and Pinkerton, 1955; Patten, 1978; Patten and Auble, 1981; Ulanowicz, 1986), dating back at least to Darwin’s *Origin of Species* in his famous pondering of an entangled bank (Bascompte and Jordano, 2014; Golley, 1993). The application of network theory has provided a formal, mathematical framework to approach systems (Bascompte and Jordano, 2014; Proulx et al., 2005) and led to the development of network ecology (Borrett et al., 2014; Patten and Witkamp, 1967; Poisot et al., 2016b).

Network ecology can be defined as the use of network models and analyses to investigate the structure, function, and evolution of ecological systems at many scales and levels of organization (Borrett et al., 2012; Eklöf et al., 2012). The influx of network thinking throughout ecology, and ecology’s contribution to the development of network science highlights the assertion that “networks are everywhere” (Lima, 2011). And, as one would expect, the field has grown rapidly, from 1% of the primary ecological literature in 1991 to over 6% in 2017 (Fig. 1A). Some examples include: applying network theory to population dynamics and spread of infectious diseases (May, 2006); description and analysis of networks of proteins in adult organisms (Stumpf et al., 2007) or during development (Hollenberg, 2007); expanding classical food webs to include parasites and non-trophic interactions (Ings et al., 2009; Kéfi et al., 2012); investigating animal movement patterns (Lédée et al., 2016) and the spatial structure of metapopulations (Dubois et al., 2016; Holstein et al., 2014); connecting biodiversity to ecosystem functioning (Creamer et al., 2016); identifying keystone species (Borrett, 2013; Zhao et al., 2016); and using social network theory in studies of animal behavior (Croft et al., 2004; Fletcher et al., 2013; Krause et al., 2003; Sih et al., 2009). Further, ideas and concepts from network ecology are being applied to investigate the sustainability of urban and industrial systems (Fang et al.,

2014; Layton et al., 2016; Xia et al., 2016) and elements of the food-energy-water nexus (Wang and Chen, 2016; Yang and Chen, 2016).

Over the past 15 years, re-occurring themes for moving network ecology forward have emerged from reviews, perspectives, and syntheses (e.g. Bascompte, 2010; Borrett et al., 2014; Poisot et al., 2015; Proulx et al., 2005). In this paper, we examine areas where the network approach is being applied to address important ecological questions and identify both challenges and opportunities for advancing the field. Among these are the need for shifting the focus toward mechanisms rather than observations, and increasing the resolution (e.g. individuals or traits as nodes and weighted edges of different interaction types) and replication of network models across different ecosystems and time (Ings et al., 2009; Poisot et al., 2016b; Woodward et al., 2010). After a brief primer of key concepts from network ecology, we discuss the following topics as they relate to these issues: the proliferation of terminology for ecological metrics with the increasing application of network methods; fully exploring the underlying assumptions of models of mechanistic processes for generating network structure; and the need for improved sharing and reproducibility of ecological network research and models. Although these topics are not new, the combination of the influx of metrics and theory and rapid increases in the computational intensity of ecology are creating novel challenges. With respect to these issues, we discuss recent advances that should be explored as tools to aid in a more effective integration of network methods for synthesis across ecological (sub)disciplines.

## 2 A primer of ecological networks: models and metrics

Prior to the introduction of network methods in ecology, the primary way of studying interactions was limited to detailed studies of behaviors and traits of individual species important to interactions, or of relationships between tightly interacting pairs of species (Carmel et al., 2013). Some ecologists were advancing whole-system methods (Lindeman, 1942; Odum, 1957); however, quantifying interactions is costly, as compared to surveys of species abundances. This has created a significant barrier to studying interactions at the scale of entire communities, either at the scale of individuals or species pairs, because the number of interactions becomes intractable. For instance, even if one assumes that only pairwise interactions occur among  $S$  species, the number of possible pairs is  $S(S-1)/2$ . Local assemblages of macrobes often have  $10^1 - 10^2$  species, and microbial diversity can easily exceed  $10^3$  OTUs (Operational Taxonomic Units).

100 This complexity of ecological systems is one reason there is a long tradition in  
 101 community ecology of studying interactions within small subsets of closely-related  
 102 species (e.g. trophic guilds) and using dimensionality reducing methods based on  
 103 multivariate, correlative approaches (Legendre et al., 2012). While some approaches  
 104 to studying subsets of species incorporate the underlying pattern of direct and in-  
 105 direct links (e. g., modules, (sensu Holt, 1997; Holt and Hoopes, 2005), the ma-  
 106 jority do not. Such limitations repeatedly have led to calls for the application of  
 107 “network thinking” to ecological questions (e.g. Golubski et al., 2016; Ings et al.,  
 108 2009; Jacoby and Freeman, 2016; Patten and Witkamp, 1967; Proulx et al., 2005;  
 109 QUINTESENCE Consortium et al., 2016; Urban and Keitt, 2001). There are now  
 110 many resources for learning about network ecology and network theory in general,  
 111 and we point the reader in the direction of excellent reviews in this area (Bascompte  
 112 and Jordano, 2007; Borrett et al., 2012; Brandes et al., 2013; Ings et al., 2009; Proulx  
 113 et al., 2005) and more comprehensive introductions (Brandes et al., 2005; Estrada,  
 114 2015; Newman, 2010).

115 Network ecology employs network theory to quantify the structure of ecological  
 116 interactions. All networks consist of sets of interacting nodes (e.g. species, non-  
 117 living nutrient pools, habitat patches) whose relationships are represented by edges  
 118 (e.g. nutrient or energy transfers, pollination, movement of individuals). Conceptu-  
 119 ally, a network is a set of things or objects with connections among them. Stated  
 120 mathematically, a network is a generic relational-model comprised of a set of objects  
 121 represented by nodes or vertices ( $N$ ) and a set of edges ( $E$ ) that map one or more  
 122 relationships among the nodes,  $G = (N, E)$ . A canonical ecological example of a net-  
 123 work is a food-web diagram, in which the nodes represent species, groups of species,  
 124 or non-living resources, and the *edges* map the relationship who-*eats*-whom.

125 The analysis of networks is inherently hierarchical, ranging from the entire net-  
 126 work down to individual nodes and edges. Depending on the characteristics and level  
 127 of detail of the information provided for a given model, there is a large number of  
 128 network analyses and metrics that can be used to characterize the system at multiple  
 129 levels (similar to Hines and Borrett, 2014; Wasserman and Faust, 1994), including:  
 130 (1) the whole network level (i.e., the entire network), (2) the sub-network level (i.e.,  
 131 groups of two or more nodes and their edges), and (3) the individual node or edge  
 132 level (Fig. 2).

133 Network-level metrics integrate information over the entire set of nodes and edges.  
 134 For example, the number of nodes (e.g., the species richness of a food web) and  
 135 the density of connections or connectance are both network-level statistics used to  
 136 describes the overall complexity of a network and have been investigated by ecologists  
 137 for over 40 years (Allesina and Tang, 2012; May, 1972).

Sub-network level analyses focus on identifying specific subsets of nodes and edges. There are a variety of groups that have different names (e.g., module, motif, cluster, clique, environ) and different methods for measurement. Sub-networks often represent more tractable and meaningful units of study than individual nodes and edges on the one hand or entire networks on the other. For example, in landscape and population ecology, the preferential movement of individuals and genes (edges) between habitat patches (nodes) has implications for conservation of populations and the design of preserves (Calabrese and Fagan, 2004; Fletcher et al., 2013; Holt and Hoopes, 2005). Also, both nodes and edges can be divided into classes. An example of this is the bipartite graph, in which interactions occur primarily between, rather than within, each class or “part” of the community. A bipartite network has only two classes of nodes, such as in a pollination network in which the community is divided into plants being pollinated and insects that do the pollination (Petanidou et al., 2008). In this network, edges representing pollination visits can only map between two nodes in the different classes.

Metrics at the individual node or edge level quantify differences in relative importance. Whether we are interested in an individual or species that transmits disease, species whose removal will result in secondary extinctions, or key habitat patches that connect fragmented landscapes, identifying important nodes is a critical component of network analysis. Another type of node or edge-level metric classifies nodes or edges according to their roles within a network. This classification can use information from differing levels. Additionally, nodes and edges can have variable characteristics. Edges can be weighted and they can map a directed relationship (as opposed to a symmetric or undirected relationship). For example, in ecosystem networks, the edges show the directed movement of energy or nutrients from one node to another by some process like feeding, and the edge weight can indicate the amount of energy or mass in the transaction (Baird and Ulanowicz, 1989; Dame and Patten, 1981). Nodes also can be weighted (e.g. size of individual, population size, biomass of a given species). Lastly, network models are flexible enough to accommodate variation in edge types and relationships among edges (e.g. hypergraphs), but analysis of these more complicated models is challenging and has only begun to be applied in ecology (e.g. Golubski et al., 2016).

### 3 Resolving network metrics

The application of network theory defines an explicit mathematical formalism that provides a potentially unifying set of terms for ecology and its inter-disciplinary applications (QUINTESSANCE Consortium et al., 2016). Ironically, the develop-

174 ment of ecological network metrics has had an opposing affect. One reason for this  
 175 is that introductions have occurred in multiple sub-disciplinary branches (Fig. 1B)  
 176 (Blüthgen, 2010; Borrett et al., 2014; Carmel et al., 2013). Having separate research  
 177 trajectories can facilitate rapid development of ideas and the process of integration  
 178 can lead to novel insights (Hodges, 2008). At the same time, these innovations in  
 179 network ecology have come at the cost of the “rediscovery” of the same network met-  
 180 rics and subsequent description of them with new terms. This has led to different  
 181 metrics with similar purposes existing in separate areas of ecology (Table 1).

182 Ecological studies using network approaches draw from a deep well of general net-  
 183 work theory (Newman, 2003, 2006; Strogatz, 2001). Ecologists broadly use network  
 184 concepts, techniques, and tools to: (1) characterize the system organization (Borrett,  
 185 2013; Croft et al., 2004; Ulanowicz, 1986); (2) investigate the consequences of the  
 186 network organization (Borrett et al., 2006; Dunne et al., 2002; Grilli et al., 2016); and  
 187 (3) identify the processes or mechanisms that might generate the observed patterns  
 188 (Allesina and Pascual, 2008; Fath et al., 2007; Guimarães et al., 2007; Poisot et al.,  
 189 2016*b*; Ulanowicz et al., 2014; Williams and Martinez, 2000). The unnecessary pro-  
 190 liferation of network metrics is exemplified by “connectance” ( $C$ ), which is used by  
 191 food-web ecologists to mean the ratio of the number of edges in the network divided  
 192 by the total number of possible edges. Elsewhere in the network science literature,  
 193 this measurement is referred to as network density (Newman et al., 2001). As an-  
 194 other example, what ecosystem ecologists have described as “average path length”  
 195 (total system through-flow divided by the total system input) (Finn, 1976) also has  
 196 been called network aggradation (Jørgensen et al., 2000). In economics, average path  
 197 length is known as the multiplier effect (Samuelson, 1948).

198 Another kind of redundancy is the creation and use of multiple statistics that  
 199 measure the same or very similar network aspects. A clear example of this is inher-  
 200 ent in the proliferation of centrality measures to indicate node or edge importance.  
 201 Network scientists have shown that many centrality metrics are correlated (Jordán  
 202 et al., 2007; Newman, 2006; Valente et al., 2008). Likewise, Borrett and Osidele  
 203 (2007) found that nine commonly reported ecosystem network analysis metrics co-  
 204 varied in 90 plausible parameterizations of a model of phosphorus biogeochemical  
 205 cycling for Lake Lanier, GA, but that all these metrics were associated strongly with  
 206 only two underlying factors. However, even a perfect correlation does not mean  
 207 that two metrics have identical properties, and they still may diverge in different  
 208 models. Therefore, it is important to have mathematically based comparisons of  
 209 metrics (Borgatti and Everett, 2006; Borrett, 2013; Kazanci and Ma, 2015; Ludovisi  
 210 and Scharler, 2017). It is incumbent on network ecologists to establish clearly the  
 211 independence and uniqueness of the descriptive metrics used.

212 From the perspective of the broader field of ecology, the proliferation of con-  
213 cepts, terms, and metrics is not a new issue (e.g. Ellison et al., 2005; Tansley, 1935).  
214 Ecologists have a long history of using network concepts and related models in mul-  
215 tiple subdomains (e.g, metapopulations, matrix population models, community co-  
216 occurrence models, ecosystems) without fully recognizing or capitalizing on the sim-  
217 ilarities of the underlying models. Each subdomain has constructed its own concepts  
218 and methods (occasionally borrowing from other areas), and established its own jar-  
219 gon that impedes scientific development. Previous suggestions for solving this issue  
220 have focused on maintaining an historical perspective of ecology (Graham and Day-  
221 ton, 2002); Blüthgen et al. (2008) is an excellent example of how this can be done  
222 through peer-reviewed literature.

223 One possible approach that would go beyond such a diffuse, literature-centered  
224 approach would be to develop a formal ontology of concepts and metrics. An on-  
225 tology is a a set of related terms that are formally defined and supported by as-  
226 sertions (Bard and Rhee, 2004). An ontology therefore provides a framework for  
227 developing concepts within a discipline and presents the opportunity for more ef-  
228 ficient synthesis across disciplinary boundaries. The concept of an ontology is not  
229 new, but more rapid sharing of ontologies and their collaborative development have  
230 been enabled by the Internet. For example, the Open Biological Ontologies (OBO,  
231 <http://www.obofoundry.org>) supports the creation and sharing of ontologies over  
232 the web. Currently, there is no OBO for a “network ecology metric” ontology, and  
233 as far as we are aware, ontologies have yet to be explored or developed for network  
234 metrics.

235 The OBO could provide a platform for harmonizing ecological network metrics,  
236 terms, and concepts. Key obstacles to such harmonization include a requirement that  
237 network ecologists work within a common framework, and the need for an individual  
238 or leadership team to periodically curate the ontology based on new developments in  
239 the field. In determining the best course of action, network ecologists could follow the  
240 example of how similar OBO projects have been managed in the past. The *FOODON*  
241 food role ontology project (<http://www.obofoundry.org/ontology/foodon.html>)  
242 contains information about “materials in natural ecosystems and food webs as well  
243 as human-centric categorization and handling of food.” It could serve as an example  
244 or even the basis of a ecological network metric ontology.



## 4 Benchmarking: Trusting our models of mechanisms

Inferences about processes in ecological systems have relied in part on the application of simulation models that generate matrices with predictable properties. As discussed in the previous section, the proliferation of network metrics points to the need for the investigation and comparison of how these metrics will behave in the context of different modeling algorithms. Once a metric or algorithm has been chosen, it is tempting apply them widely to empirical systems to detect patterns, but before research proceeds, a process of “benchmarking” with artificial matrices that have predefined amounts of structure and randomness should be used to examine the behavior of the algorithms and the metrics that are applied to them.

Benchmarking of ecological models developed from null model analysis in community ecology (Atmar and Patterson, 1993; Connor and Simberloff, 1979; Gotelli and Ulrich, 2012). Null models are specific examples of randomization or Monte Carlo tests (Manly, 2007) that estimate a frequentist  $P$  value, the tail probability of obtaining the value of some metric if the null hypothesis were true (Gotelli and Graves, 1996). The aim of a null model is to determine if the structure of an observed ecological pattern in space or time is incongruous with what would be expected given the absence of a causal mechanism. A metric of structure calculated for a single empirical data set is compared to the distribution of the same metric calculated for a collection of a large number of randomizations of the empirical data set. The data are typically randomized by reshuffling some elements while holding other elements constant to incorporate realistic constraints. Comparison with a suite of null models in which different constraints are systematically imposed or relaxed may provide a better understanding of the factors that contribute most to patterns in the network (see Box 1). However, the devil remains in the details and there are also a variety of ways to randomize data and impose constraints to construct a useful null model. If the null model is too simplistic, such as a model in which edges and nodes are sampled with uniform probability, it will always be rejected and provide little insight into important ecological patterns, regardless of what metric is used. At the other extreme, if the null model incorporates too many constraints from the data, it will be difficult or impossible to reject the null hypothesis, even though the network may contain interesting structure.

In network theory, the Erdos-Renyi (ER, (Erdős and Rényi, 1959)) model is a now-classic example of a model used to generate networks via a random process for creating matrix structure. The ER model is a random graph that starts with an  $N \times N$  adjacency matrix of nodes and assigns to it  $K$  edges between randomly chosen

282 pairs of nodes. The ER model has been applied in ecology to address questions about  
 283 the relationship between stability and complexity (May, 1972) and the structure of  
 284 genetic networks (Kauffman et al., 2003). For example, randomized networks have  
 285 been used to link motifs (Milo et al., 2002) to network assembly (Baiser et al., 2016),  
 286 stability (Allesina and Pascual, 2008; Borrelli et al., 2015), and persistence in food  
 287 webs (Stouffer and Bascompte, 2010).

288 In addition to the random matrix approaches of null and ER models, there are  
 289 other, more complex algorithms that are used to generate structured matrices. Per-  
 290 haps one of the best known in network theory is the Barabasi-Albert (BA, Barabási  
 291 and Albert 1999) model, which adds nodes and edges to a growing network with  
 292 a greater probability of adding edges to nodes with a higher degree. The BA algo-  
 293 rithm is similar to ecological network algorithms that generate non-random structure,  
 294 because of either direct influence or similar processes operating in systems of inter-  
 295 est. Some of these models include processes of “preferential attachment” in which  
 296 organisms tend to interact with the same, common species. Food-web modeling al-  
 297 gorithms also have been developed that use a trait-based approach (e.g. Allesina and  
 298 Pascual, 2009), consumer-resource models (Yodzis and Innes, 1992), niches (Williams  
 299 and Martinez, 2000), cyber-ecosystem algorithms (Fath, 2004), and cascade models  
 300 (Allesina and Pascual, 2009; Allesina and Tang, 2012; Cohen and Luczak, 1992).

301 The statistical behavior of some models and metrics can be understood ana-  
 302 lytically. For example, the networks generated by the BA algorithm display degree  
 303 distributions that approximate a power-law distribution, like many real-world “scale-  
 304 free” networks (Albert et al., 2002). If the network is sparse (i.e.  $(K \ll N^2)$ ), the  
 305 degree distribution of the network should follow a Poisson distribution. However, as  
 306 new models and metrics are introduced, new benchmarking should be done and com-  
 307 pared to previous results. Newman et al. (2016) is one example of how benchmarking  
 308 can be used for investigating processes operating on ecological networks. Ludovisi  
 309 and Scharler (2017) advocate the same approach for the analysis of network models  
 310 in general. The **benchmark** (Eugster and Leisch, 2008) package in R (R Core Team,  
 311 2017) is a general algorithm-testing software package that provides a useful starting  
 312 point.

## 313 5 Reproducibility: Open-data, Open-source and 314 Provenance

315 As analyses of network models increase in computational intensity, there is a concomi-  
 316 tant increase in the need for new tools to track and share key computational details.

317 This need is compounded when models incorporate data from multiple sources or  
318 analyses involve random processes. The combination of the volume of data and com-  
319 putational intensity of studies of ecological networks further increases the burden on  
320 ecologists to provide information needed to adequately reproduce datasets, analyses,  
321 and results. As the sharing and reproducibility of scientific studies are both essential  
322 for advances to have lasting impact, finding easier, faster, and generally more conve-  
323 nient ways to record and report relevant information for ecological network studies  
324 is imperative for advancing the field.

325 Sharing data and open-source code have become established in ecology, and net-  
326 work ecologists are now producing more network models and data (e.g. Fig. 1A).  
327 These include not only ecological interaction networks, but also an influx of other rele-  
328 vant networks, including ecological genomic networks generated by next-generation,  
329 high-throughput sequencing technologies (Langfelder and Horvath, 2008; Zinkgraf  
330 et al., 2017). There are now multiple web-accessible scientific databases (e.g. NCBI,  
331 Data Dryad, Dataverse) and at least four databases have been constructed specifically  
332 to curate ecological network data: including “Kelpforest” (Beas-Luna et al., 2014),  
333 “The Web of Life” (Fortuna et al., 2014), “Mangal” ecological network database  
334 (Poisot et al., 2015) and the “Interaction Web Database” ([https://www.nceas.  
335 ucsb.edu/interactionweb/resources.html](https://www.nceas.ucsb.edu/interactionweb/resources.html)).

336 The increase in ecological network data is linked to an increasing rate of shared  
337 analytical code and other open-source software. It is now commonplace for ecologists  
338 to have a working knowledge of one or more programming languages, such as R,  
339 Python, SAS, MatLab, Mathematica, or SPSS. Multiple software packages exist for  
340 doing ecological analyses, including ecological network analyses. In addition to the  
341 general network analysis packages available in R, there are at least two packages  
342 aimed specifically at ecological network analysis: **bipartite** and **enaR**. The former  
343 provides functions drawn largely from community ecology (Dormann et al., 2009),  
344 whereas the latter provides a suite of algorithms developed in the ecosystem network  
345 analysis literature (Borrett and Lau, 2014; Lau et al., 2015).

346 Although, ecology has long had a culture of keeping records of important re-  
347 search details, such as field and lab notebooks, these practices put all of the burden  
348 of recording “metadata” on the researcher. Manual record-keeping methods, even  
349 when conforming to metadata standards (Boose et al., 2007, e.g. EML, see), do not  
350 take advantage of the power of the computational environment. Data-provenance  
351 methods aim to provide a means to collect formalized information about computa-  
352 tional processes, ideally in a way that aids the reproducibility of studies with minimal  
353 impact on the day-to-day activities of researchers (Boose et al., 2007). These tech-  
354 niques have been applied in other areas of research and could provide an effective

means for documenting the source and processing of data from the raw state into a model (Boose and Lerner, 2017).

The reproducibility of scientific studies is imperative for advances to have lasting impact through the independent verification of results. Although this has been an ongoing topic of discussion in ecology (Ellison, 2010; Parker et al., 2016), the need was highlighted by a recent survey finding issues with reproduction of studies across many scientific disciplines (Baker, 2016). There is significant motivation from within the ecological community to move toward providing detailed information about computational workflows for both repeatability and reproducibility, which includes repetition by the original investigator (Lowndes et al., 2017). It is also important in network ecology for data sources and methods for model construction be standardized and transparent, and that models be curated and shared (McNutt et al., 2016).

Collecting details, such as those enabled by data-provenance capture software, is one innovative way forward. These tools have been developing in the computer-science domain for decades; however, only recently have they gained a foothold in ecology (Boose et al., 2007; Ellison, 2010) or the broader scientific community. Although there are many challenges in the development and application of data-provenance principles, multiple software packages do exist for collecting data provenance in the context of scientific investigations. Two provenance capture packages exist in R, the **recordr** package associated with the DataOne repository (Cao et al., 2016) and **RDataTracker** (Lerner and Boose, 2014). In addition, although they do not collect formal data provenance, there are methods developed for “literate computing” that help to collect code along with details about the code and the intention of the analyses (e.g., the Jupyter notebook project: (Shen and Barabasi, 2014)).

For ecological networks, there is software that captures the “data pedigree” of food-web models, but it does not capture data provenance. Data pedigree was initially implemented in the EcoPath food-web modeling package (Guesnet et al., 2015; Heymans et al., 2016) to define confidence intervals and precision estimates for network edges. It has been developed further to allow for the use of informative priors in Bayesian modeling of ecological networks. This is done by linking models to the literature sources from which estimates were derived, an approach that is similar to incorporating metadata information within databases of ecological networks. Although this approach focuses only on a subcomponent of provenance, this still is a promising way to address the issue that networks, network metrics, and simulation models used to analyze them commonly assume a lack of uncertainty (*cf.* Borrett and Osidele, 2007; Kauffman et al., 2003; Kones et al., 2009), and typically ignore inaccuracy in the empirical data (Ascough et al., 2008; Gregr and Chan, 2014).

## 6 Moving Forward

Development and application of new technologies (e.g. sequencing methods and computational, data-driven approaches) have the potential to increase both the abundance and quality of ecological networks. For the future development of network ecology, there is a pressing need not only to share data and code, but also to integrate and use the large amounts of information enabled by technological advances. For example, synthetic networks (i.e. networks merging models from different studies, and *sensu* Poisot et al., 2016a) are a promising new direction; however, the structural properties of synthetic networks and the behavior of network metrics applied to them will require careful investigation, including the application of systematic benchmarking. Multi-trophic networks provide a precedence for these studies to move forward, but synthesizing models from across many different sources produces new challenges for developing and benchmarking metrics, as well as an opportunity for new technologies, like data provenance, to help establish better connections among studies and researchers.

The burgeoning of “open” culture in the sciences (Hampton et al., 2014) also has the potential to serve as a resource for models and a clearinghouse for resolving the validity of metrics, models, and algorithms. First, because code is openly shared, functions used to calculate metrics are open for inspection and, if coded and documented clearly using software “best-practices” (e.g. Noble, 2009; Visser et al., 2015), the code provides a transparent documentation of how a metric is implemented and its computational similarity to other metrics. Second, enabled by the ability to write their own functions and code, researchers can do numerical investigations of the similarities among metrics. Through comparison of metrics calculated on the same or similar network models, a researcher could at least argue, for a given set of models, that two or more metrics produce similar results. Third, data provenance provides a useful tool to aide in the dissemination and synthesis of network models and increases the reproducibility of ecological network studies, including those documenting new metrics and benchmarking those metrics and associated algorithms for generating or analyzing empirical models. Last, as with data provenance, formalizing ecological network metrics and concepts requires a mathematically rigorous foundation that is developed by the community of researchers working along parallel lines of inquiry. Whether this is done through an ontological approach or some other formalized “clearing-house,” an open process of exchange that integrates multiple perspectives is essential to prevent the rapid dilution of concepts in ecological network research as these concepts continue to proliferate, develop and evolve.

Over half a century ago, Robert MacArthur published his first paper on the rela-

430 tionship between diversity and stability, initiating multiple research trajectories that  
 431 have now become the mainstay of many ecological research programs (MacArthur,  
 432 1955). The theory that MacArthur applied was based on flows of energy through  
 433 networks of interacting species. Thus, network theory is at the roots of one of the  
 434 most widely studied topics in ecology and is now a part of the broader context of  
 435 integration across many scientific disciplines that is aimed at consilience of theory  
 436 (Wilson, 1999). The synthesis of ecological concepts through the mathematically  
 437 rigorous “lingua franca” of network terminology has the potential to unify theories  
 438 across disciplines. As with previous concepts (e.g. keystone species, foundation  
 439 species, ecosystem engineer), greater clarity and less redundancy will come about  
 440 as network methods are used more commonly and researchers compare the mathe-  
 441 matical and computational underpinnings of the metrics that they are using. With  
 442 the increased use of these approaches, the network concept has and will continue to  
 443 serve as a common model that transcends disciplines and has the potential to serve  
 444 as an inroad for new approaches. With thoughtful dialogue across sub-disciplines  
 445 and among research groups, further infusion of network theory and methods will  
 446 continue to advance ecology.

## 447 References

- 448 Albert, R., A. L. Barabasi, and A.-L. Barabási. 2002. Statistical mechan-  
 449 ics of complex networks. *Reviews of Modern Physics* **74**:47–97. URL <http://journals.aps.org/rmp/abstract/10.1103/RevModPhys.74.47>  
 450 <http://dx.doi.org/10.1103/RevModPhys.74.47>  
 451 <http://dx.doi.org/10.1103/RevModPhys.74.47>  
 452 <http://dx.doi.org/10.1103/RevModPhys.74.47>.
- 453 Allesina, S., and M. Pascual. 2008. Network structure, predatorprey modules, and  
 454 stability in large food webs. *Theoretical Ecology* **1**:55–64. URL <http://www.springerlink.com/index/10.1007/s12080-007-0007-8>.
- 456 Allesina, S., and M. Pascual. 2009. Food web models: A plea for groups. *Ecology*  
 457 *Letters* **12**:652–662.
- 458 Allesina, S., and S. Tang. 2012. Stability criteria for complex ecosystems. *Nature*  
 459 **483**:205–8. URL <http://dx.doi.org/10.1038/nature10832>.
- 460 Ascough, J., H. Maier, J. Ravalico, and M. Strudley. 2008. Future research chal-  
 461 lenges for incorporation of uncertainty in environmental and ecological decision-

- 462 making. *Ecological Modelling* **219**:383–399. URL <http://www.sciencedirect.com/science/article/pii/S0304380008003554>.  
463
- 464 Atmar, W., and B. D. Patterson. 1993. The measure of order and disorder in  
465 the distribution of species in fragmented habitat. *Oecologia* **96**:373–382. URL  
466 <http://link.springer.com/10.1007/BF00317508>.
- 467 Baird, D., and R. E. Ulanowicz. 1989. The seasonal dynamics of the Chesapeake  
468 Bay ecosystem. *Ecological Monographs* **59**:329–364.
- 469 Baiser, B., R. Elhesha, and T. Kahveci. 2016. Motifs in the assembly of food web  
470 networks. *Oikos* **125**:480–491. URL <http://dx.doi.org/10.1111/oik.02532>.
- 471 Baiser, B., J. L. Lockwood, D. La Puma, and M. F. J. Aronson. 2008. A per-  
472 fect storm: two ecosystem engineers interact to degrade deciduous forests of New  
473 Jersey. *Biological Invasions* **10**:785–795. URL <http://link.springer.com/10.1007/s10530-008-9247-9>.  
474
- 475 Baker, M. 2016. 1,500 scientists lift the lid on reproducibility. *Nature* **533**:452–454.  
476 URL <http://www.nature.com/doifinder/10.1038/533452a>.
- 477 Barabási, A.-L., and R. Albert. 1999. Emergence of scaling in random networks.  
478 *Science* **286**:509–512.
- 479 Barabási, A.-L., R. Albert, and H. Jeong. 2000. Scale-free characteristics of random  
480 networks: the topology of the world-wide web. *Physica A: statistical mechanics*  
481 *and its applications* **281**:69–77.
- 482 Bard, J. B. L., and S. Y. Rhee. 2004. Ontologies in biology: design, applications  
483 and future challenges. *Nature Reviews Genetics* **5**:213–222. URL <http://www.nature.com/doifinder/10.1038/nrg1295>.  
484
- 485 Bascompte, J. 2010. Ecology. Structure and dynamics of ecological networks. *Science*  
486 (New York, N.Y.) **329**:765–6. URL <http://www.sciencemag.org/content/329/5993/765.short>.  
487
- 488 Bascompte, J., and P. Jordano. 2007. Plant-Animal Mutualistic Networks: The Ar-  
489 chitecture of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*  
490 **38**:567–593.
- 491 Bascompte, J., and P. Jordano. 2014. Mutualistic networks. Princeton University  
492 Press.

- 493 Beas-Luna, R., M. Novak, M. H. Carr, M. T. Tinker, A. Black, J. E. Caselle,  
494 M. Hoban, D. Malone, and A. Iles. 2014. An online database for informing  
495 ecological network models: <http://kelpforest.ucsc.edu>. PloS one **9**:e109356. URL  
496 <http://dx.plos.org/10.1371/journal.pone.0109356>.
- 497 Blüthgen, N. 2010. Why network analysis is often disconnected from community ecol-  
498 ogy: A critique and an ecologist's guide. Basic and Applied Ecology **11**:185–195.  
499 URL <http://linkinghub.elsevier.com/retrieve/pii/S1439179110000125>.
- 500 Blüthgen, N., J. Fründ, D. P. Vázquez, and F. Menzel. 2008. WHAT DO INTERAC-  
501 TION NETWORK METRICS TELL US ABOUT SPECIALIZATION AND BI-  
502 OLOGICAL TRAITS. Ecology **89**:3387–3399. URL <http://www.esajournals.org/doi/abs/10.1890/07-2121.1>.
- 503
- 504 Bonacich, P., 1987. Power and Centrality: A Family of Measures.
- 505 Boose, E. R., A. M. Ellison, L. J. Osterweil, L. a. Clarke, R. Podorozhny, J. L. Hadley,  
506 A. Wise, and D. R. Foster. 2007. Ensuring reliable datasets for environmental  
507 models and forecasts. Ecological Informatics **2**:237–247.
- 508 Boose, E. R., and B. S. Lerner, 2017. Replication of data analyses: provenance in R.
- 509 Borgatti, S. P., and M. G. Everett. 2006. A Graph-theoretic perspective on centrality.  
510 Social Networks **28**:466–484. URL [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/pii/S0378873305000833)  
511 [article/pii/S0378873305000833](http://www.sciencedirect.com/science/article/pii/S0378873305000833).
- 512 Borrelli, J. J., S. Allesina, P. Amarasekare, R. Arditi, I. Chase, J. Damuth, R. D.  
513 Holt, D. O. Logofet, M. Novak, R. P. Rohr, A. G. Rossberg, M. Spencer, J. K.  
514 Tran, and L. R. Ginzburg. 2015. Selection on stability across ecological scales.  
515 Trends in Ecology & Evolution **30**:417–425.
- 516 Borrett, S. R. 2013. Throughflow centrality is a global indicator of the functional  
517 importance of species in ecosystems. Ecological Indicators **32**:182–196. URL <http://dx.doi.org/10.1016/j.ecolind.2013.03.014>.  
518
- 519 Borrett, S. R., W. Bridewell, P. Langely, and K. R. Arrigo. 2006. A method for  
520 representing and developing process models. Ecological Complexity **4**:28. URL  
521 <http://arxiv.org/abs/q-bio/0605025>.
- 522 Borrett, S. R., R. R. Christian, and R. E. Ulanowicz, 2012. Network Ecol-  
523 ogy (Revised). Pages 1767–1772 in A. El-Shaarawi and W. Piegorsch, editors.



524 Encyclopedia of Environmetrics (2nd edition). John Wiley and Sons, Chinch-  
525 ester, second edition. URL [http://people.uncw.edu/borretts/documents/](http://people.uncw.edu/borretts/documents/BorrettChristianUlanowicz2012NetworkEcology.pdf)  
526 [BorrettChristianUlanowicz2012NetworkEcology.pdf](http://people.uncw.edu/borretts/documents/BorrettChristianUlanowicz2012NetworkEcology.pdf).

527 Borrett, S. R., B. D. Fath, and B. C. Patten. 2007. Functional integration of  
528 ecological networks through pathway proliferation. *J. Theor. Biol.* **245**:98–111.

529 Borrett, S. R., and M. K. Lau. 2014. {enaR}: An {R} package for Ecosystem  
530 Network Analysis. *Methods in Ecology and Evolution* **11**:1206–1213.

531 Borrett, S. R., J. Moody, and A. Edelman. 2014. The rise of Network Ecology: Maps  
532 of the topic diversity and scientific collaboration. *Ecological Modelling* page 18.  
533 URL <http://arxiv.org/abs/1311.1785>.

534 Borrett, S. R., and O. O. Osidele. 2007. Environ indicator sensitivity to flux un-  
535 certainty in a phosphorus model of Lake Sidney Lanier, USA. *Ecological Mod-*  
536 *elling* **200**:371–383. URL [http://www.sciencedirect.com/science/article/](http://www.sciencedirect.com/science/article/B6VBS-4M57HDN-2/2/eeeeed79df4b4a4d4787c2ce6f3089af6)  
537 [B6VBS-4M57HDN-2/2/eeeeed79df4b4a4d4787c2ce6f3089af6](http://www.sciencedirect.com/science/article/B6VBS-4M57HDN-2/2/eeeeed79df4b4a4d4787c2ce6f3089af6).

538 Borrvall, C., B. Ebenman, and T. Jonsson, 2000. Biodiversity lessens the risk of  
539 cascading extinction in model food webs.

540 Brandes, U., T. Erlebach, and Gesellschaft fur Informatik. 2005. Network analysis :  
541 methodological foundations. Springer.

542 Brandes, U., G. Robins, A. McCranie, and S. Wasserman. 2013. What is network  
543 science? *Network Science* **1**:1–15.

544 Calabrese, J. M., and W. F. Fagan. 2004. A comparison-shopper’s guide to connec-  
545 tivity metrics. *Frontiers in Ecology and the Environment* **2**:529–536. URL [http://doi.wiley.com/10.1890/1540-9295\(2004\)002\[0529:ACGTCM\]2.0.CO;2](http://doi.wiley.com/10.1890/1540-9295(2004)002[0529:ACGTCM]2.0.CO;2).

547 Cao, Y., C. Jones, V. Cuevas-Vicenttín, M. B. Jones, B. Ludäscher, T. McPhillips,  
548 P. Missier, C. Schwalm, P. Slaughter, D. Vieglais, L. Walker, and Y. Wei,  
549 2016. DataONE: A Data Federation with Provenance Support. Pages 230–234  
550 . Springer International Publishing. URL [http://link.springer.com/10.1007/](http://link.springer.com/10.1007/978-3-319-40593-3_{\_}28)  
551 [978-3-319-40593-3\\_{\\\_}28](http://link.springer.com/10.1007/978-3-319-40593-3_{\_}28).

552 Carmel, Y., R. Kent, A. Bar-Massada, L. Blank, J. Liberzon, O. Nezer, G. Sapir, and  
553 R. Federman. 2013. Trends in ecological research during the last three decades—a  
554 systematic review. *PloS one* **8**:e59813.

- 555 Cherrett, J. M., 1989. Key concepts: The results of a survey of our members'  
556 opinions. Pages 1–16 in J. M. Cherrett, A. D. Bradshaw, F. B. Goldsmith, P. G.  
557 Grubb, and J. R. Krebs, editors. Ecological concepts: The contribution of ecology  
558 to an understanding of the natural world. Blackwell Scientific Publications, Oxford,  
559 UK.
- 560 Cohen, J. E., and T. Łuczak. 1992. Trophic levels in community food webs.  
561 Evolutionary Ecology **6**:73–89. URL [http://link.springer.com/10.1007/  
562 BF02285335](http://link.springer.com/10.1007/BF02285335).
- 563 Colwell, R. K., and D. W. Winkler. 1984. A null model for null models in biogeog-  
564 raphy. Princeton University Press.
- 565 Connor, E. F., and D. Simberloff. 1979. The Assembly of Species Communi-  
566 ties: Chance or Competition? Ecology **60**:1132. URL [http://www.jstor.org/  
567 stable/1936961?origin=crossref&GotoISI://A1979JS62200008](http://www.jstor.org/stable/1936961?origin=crossref&GotoISI://A1979JS62200008).
- 568 Council, N. R. 2003. Neon. National Academies Press, Washington, D.C. URL  
569 <http://www.nap.edu/catalog/10807>.
- 570 Creamer, R., S. Hannula, J. Van Leeuwen, D. Stone, M. Rutgers, R. Schmelz,  
571 P. De Ruiter, N. B. Hendriksen, T. Bolger, M.-L. Bouffaud, et al. 2016. Eco-  
572 logical network analysis reveals the inter-connection between soil biodiversity and  
573 ecosystem function as affected by land use across Europe. Applied Soil Ecology  
574 **97**:112–124.
- 575 Croft, D. P., J. Krause, and R. James. 2004. Social networks in the guppy  
576 (*Poecilia reticulata*). Proc. Royal Soc. Lond. B. **271**:S516–S519.
- 577 Dame, R. F., and B. C. Patten. 1981. Analysis of energy flows in an intertidal oyster  
578 reef. Marine Ecology Progress Series **5**:115–124. URL [http://www.int-res.com/  
579 articles/meps/5/m005p115.pdf](http://www.int-res.com/articles/meps/5/m005p115.pdf).
- 580 Dayton, P. K. 1972. Toward an understanding of community resilience and the  
581 potential effects of enrichment to the benthos at McMurdo Sound, Antarctica.  
582 Proceedings of the Colloquium on Conservation Problems in Antarctica pages 81–  
583 96. URL <http://daytonlab.ucsd.edu/Publications/Pubs.htm>.
- 584 Dormann, C. F., C. F. Dormann, J. Fründ, N. Blüthgen, and B. Gruber. 2009.  
585 Indices, Graphs and Null Models: Analyzing Bipartite Ecological Networks.  
586 Open Ecology Journal **2**:7–24. URL [http://citeseerx.ist.psu.edu/viewdoc/  
587 summary?doi=10.1.1.323.9930](http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.323.9930).

- 588 Dubois, M., V. Rossi, E. Ser-Giacomi, S. Arnaud-Haond, C. López, and  
589 E. Hernández-García. 2016. Linking basin-scale connectivity, oceanography and  
590 population dynamics for the conservation and management of marine ecosystems.  
591 *Global Ecology and Biogeography* .
- 592 Dunne, J. a., R. J. Williams, and N. D. Martinez. 2002. Network structure and  
593 biodiversity loss in food webs: Robustness increases with connectance. *Ecology*  
594 *Letters* **5**:558–567.
- 595 Eklöf, A., and B. Ebenman. 2006. Species loss and secondary extinctions in simple  
596 and complex model communities. *Journal of Animal Ecology* **75**:239–246.
- 597 Eklöf, A., M. R. Helmus, M. Moore, and S. Allesina. 2012. Relevance of evolutionary  
598 history for food web structure. *Proceedings. Biological sciences / The Royal So-*  
599 *cietty* **279**:1588–96. URL [http://rspb.royalsocietypublishing.org/content/](http://rspb.royalsocietypublishing.org/content/early/2011/11/10/rspb.2011.2149.full)  
600 [early/2011/11/10/rspb.2011.2149.full](http://rspb.royalsocietypublishing.org/content/early/2011/11/10/rspb.2011.2149.full).
- 601 Ellison, A. M. 2010. Repeatability and transparency in ecological research. *Ecology*  
602 **91**:2536–2539.
- 603 Ellison, A. M., M. S. Bank, B. D. Clinton, E. A. Colburn, K. Elliott, C. R. Ford, D. R.  
604 Foster, B. D. Kloeppel, J. D. Knoepp, G. M. Lovett, J. Mohan, D. A. Orwig, N. L.  
605 Rodenhouse, W. V. Sobczak, K. A. Stinson, J. K. Stone, C. M. Swan, J. Thompson,  
606 B. Von Holle, and J. R. Webster. 2005. Loss of foundation species: consequences  
607 for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and*  
608 *the Environment* **3**:479–486. URL [http://www.esajournals.org/doi/abs/10.](http://www.esajournals.org/doi/abs/10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2?journalCode=fron)  
609 [1890/1540-9295\(2005\)003\[0479:LOFSCF\]2.0.CO;2?journalCode=fron](http://www.esajournals.org/doi/abs/10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2?journalCode=fron).
- 610 Erdős, P., and a. Rényi. 1959. On random graphs. *Publicationes Mathematicae*  
611 **6**:290–297.
- 612 Estrada, E., 2015. *Introduction to Complex Networks: Structure and Dynamics*.  
613 Pages 93–131 . Springer International Publishing. URL [http://link.springer.](http://link.springer.com/10.1007/978-3-319-11322-7{\_}3)  
614 [com/10.1007/978-3-319-11322-7{\\\_}3](http://link.springer.com/10.1007/978-3-319-11322-7{\_}3).
- 615 Eugster, M. J. A., and F. Leisch, 2008. Bench Plot and Mixed Effects Models: First  
616 Steps toward a Comprehensive Benchmark Analysis Toolbox. Pages 299–306 *in*  
617 P. Brito, editor. *Compstat 2008—Proceedings in Computational Statistics*. Physica  
618 Verlag, Heidelberg, Germany. URL <http://epub.ub.uni-muenchen.de/3206/>.

- 619 Fang, D., B. D. Fath, B. Chen, and U. M. Scharler. 2014. Network environ  
620 analysis for socio-economic water system. *Ecological Indicators* URL [http:](http://www.sciencedirect.com/science/article/pii/S1470160X14001976)  
621 [//www.sciencedirect.com/science/article/pii/S1470160X14001976](http://www.sciencedirect.com/science/article/pii/S1470160X14001976).
- 622 Fath, B. D. 2004. Network analysis applied to large-scale cyber-ecosystems. *Ecolog-*  
623 *ical Modeling* **171**:329–337.
- 624 Fath, B. D., U. M. Scharler, R. E. Ulanowicz, and B. Hannon. 2007. Ecological  
625 network analysis: network construction. *Ecological Modelling* **208**:49–55. URL  
626 <http://www.sciencedirect.com/science/article/pii/S0304380007002517>.
- 627 Finn, J. T. 1976. Measures of ecosystem structure and function derived from analysis  
628 of flows. *Journal of theoretical biology* **56**:363–380.
- 629 Finn, J. T. 1980. Flow analysis of models of the Hubbard Brook ecosystem. *Ecology*  
630 **61**:562–571.
- 631 Fletcher, R. J., A. Revell, B. E. Reichert, W. M. Kitchens, J. D. Dixon, and J. D.  
632 Austin. 2013. Network modularity reveals critical scales for connectivity in ecol-  
633 ogy and evolution. *Nature Communications* **4**. URL [http://www.nature.com/](http://www.nature.com/doifinder/10.1038/ncomms3572)  
634 [doifinder/10.1038/ncomms3572](http://www.nature.com/doifinder/10.1038/ncomms3572).
- 635 Fortuna, M. A., R. Ortega, and J. Bascompte. 2014. The Web of Life. [www.web-of-](http://www.web-of-life.es)  
636 [life.es](http://www.web-of-life.es) URL <http://arxiv.org/abs/1403.2575>[www.web-of-life.es](http://www.web-of-life.es).
- 637 Freeman, L. C. 1979. Centrality in networks. I. Conceptual clarificaion. *Social*  
638 *Networks* **1**:215–239.
- 639 Golley, F. 1993. A history of the ecosystem concept in ecology: More than the sum  
640 of the parts. Yale University Press, New Haven, CT.
- 641 Golubski, A. J., E. E. Westlund, J. Vandermeer, and M. Pascual. 2016. Ecological  
642 Networks over the Edge: Hypergraph Trait-Mediated Indirect Interaction (TMII)  
643 Structure. *Trends in Ecology & Evolution* URL [http://www.cell.com/article/](http://www.cell.com/article/S0169534716000513/fulltext)  
644 [S0169534716000513/fulltext](http://www.cell.com/article/S0169534716000513/fulltext).
- 645 Gotelli, N. J. 2000. Null model analysis of species co-occurrence patterns. *Ecology*  
646 **81**:2606–2621.
- 647 Gotelli, N. J., R. M. Dorazio, A. M. Ellison, and G. D. Grossman. 2010. Detecting  
648 temporal trends in species assemblages with bootstrapping procedures and hierar-  
649 chical models. *Philosophical transactions of the Royal Society of London. Series*  
650 *B, Biological sciences* **365**:3621–3631.

- 651 Gotelli, N. J., A. M. Ellison, and B. A. Ballif. 2012. Environmental proteomics, biodi-  
652 versity statistics and food-web structure. *Trends in ecology & evolution* **27**:436–42.  
653 URL [/pmcc/articles/PMC3392467/?report=abstract](http://pmcc/articles/PMC3392467/?report=abstract).
- 654 Gotelli, N. J., and G. R. Graves. 1996. Null models in ecology. *Ecol-*  
655 *ogy* **14**:368. URL [http://books.google.com/books?id=vjsBAAAACAAJ{\&  
656 }printsec=frontcover](http://books.google.com/books?id=vjsBAAAACAAJ{\& }printsec=frontcover).
- 657 Gotelli, N. J., and W. Ulrich, 2012. Statistical challenges in null model analysis.
- 658 Graham, M. H., and P. K. Dayton. 2002. ON THE EVOLUTION OF  
659 ECOLOGICAL IDEAS: PARADIGMS AND SCIENTIFIC PROGRESS. *Ecol-*  
660 *ogy* **83**:1481–1489. URL [http://doi.wiley.com/10.1890/0012-9658\(2002\)  
661 083\[1481:0TE0EI\]2.0.CO;2](http://doi.wiley.com/10.1890/0012-9658(2002)083[1481:0TE0EI]2.0.CO;2).
- 662 Gregr, E. J., and K. M. A. Chan. 2014. Leaps of Faith: How Implicit Assumptions  
663 Compromise the Utility of Ecosystem Models for Decision-making. *BioScience*  
664 **65**:43–54. URL [http://bioscience.oxfordjournals.org/content/65/1/43.  
665 full](http://bioscience.oxfordjournals.org/content/65/1/43.full).
- 666 Grilli, J., T. Rogers, and S. Allesina. 2016. Modularity and stability in ecological  
667 communities. *Nature Communications* **7**.
- 668 Guesnet, V., G. Lassalle, A. Chaalali, K. Kearney, B. Saint-Béat, B. Karimi,  
669 B. Grami, S. Tecchio, N. Niquil, and J. Lobry. 2015. Incorporating food-web  
670 parameter uncertainty into Ecopath-derived ecological network indicators. *Eco-*  
671 *logical Modelling* **313**:29–40. URL [http://www.sciencedirect.com/science/  
672 article/pii/S0304380015002495](http://www.sciencedirect.com/science/article/pii/S0304380015002495).
- 673 Guimarães, P. R., G. Machado, M. A. M. de Aguiar, P. Jordano, J. Bascompte,  
674 A. Pinheiro, and S. F. Dos Reis. 2007. Build-up mechanisms determining the  
675 topology of mutualistic networks. *Journal of theoretical biology* **249**:181–9. URL  
676 <http://www.ncbi.nlm.nih.gov/pubmed/17889903>.
- 677 Hampton, S. E., S. Anderson, S. C. Bagby, and C. Gries. 2014. The Tao of Open  
678 Science for Ecology. *PeerJ PrePrints* pages 1–30. URL [https://peerj.com/  
679 preprints/549/](https://peerj.com/preprints/549/).
- 680 Heymans, J. J., M. Coll, J. S. Link, S. Mackinson, J. Steenbeek, C. Walters, and  
681 V. Christensen. 2016. Best practice in Ecopath with Ecosim food-web models for  
682 ecosystem-based management. *Ecological Modelling* **331**:173–184.

- Hines, D. E., and S. R. Borrett. 2014. A comparison of network, neighborhood, and node levels of analyses in two models of nitrogen cycling in the Cape Fear River Estuary. *Ecological Modelling* **293**:210–220. URL <http://www.sciencedirect.com/science/article/pii/S0304380013005607>.
- Hodges, K. E. 2008. Defining the problem: terminology and progress in ecology. *Frontiers in Ecology and the Environment* **6**:35–42. URL <http://doi.wiley.com/10.1890/060108>.
- Hollenberg, D. 2007. On the evolution and dynamics of biological networks. *Rivista di biologia* **100**:93–118. URL <http://www.ncbi.nlm.nih.gov/pubmed/17592821>.
- Holstein, D. M., C. B. Paris, and P. J. Mumby. 2014. Consistency and inconsistency in multispecies population network dynamics of coral reef ecosystems. *Marine Ecology Progress Series* **499**:1–18.
- Holt, R., 1997. Community modules. Pages 333–349 *in* Multitrophic interactions in terrestrial ecosystems, 36th Symposium of the British Ecological Society. Blackwell Science Oxford.
- Holt, R. D., and M. F. Hoopes. 2005. Food web dynamics in a metacommunity context. *Metacommunities: Spatial dynamics and ecological communities* pages 68–93.
- Ings, T. C., J. M. Montoya, J. Bascompte, N. Blüthgen, L. Brown, C. F. Dormann, F. Edwards, D. Figuerola, U. Jacob, J. I. Jones, R. B. Lauridsen, M. E. Ledger, H. M. Lewis, J. M. Olesen, F. J. F. van Veen, P. H. Warren, and G. Woodward. 2009. Ecological networks—beyond food webs. *The Journal of animal ecology* **78**:253–69. URL <http://www.ncbi.nlm.nih.gov/pubmed/19120606>.
- Jacoby, D. M. P., and R. Freeman. 2016. Emerging network-based tools in movement ecology. *Trends in Ecology & Evolution* **31**:301–314.
- Jones, C. G., J. H. Lawton, and M. Shachak. 1994. Organisms as Ecosystem Engineers. *Oikos* **69**:373–386.
- Jordán, F., Z. Benedek, and J. Podani. 2007. Quantifying positional importance in food webs: A comparison of centrality indices. *Ecological Modelling* **205**:270–275.
- Jørgensen, S. E., B. C. Patten, and M. Straškraba. 2000. Ecosystems emerging: 4. Growth. *Ecological Modelling* **126**:249–284.

- 714 Kauffman, S., C. Peterson, B. R. Samuelsson, C. Troein, and P. W. Ander-  
715 son. 2003. Random Boolean network models and the yeast transcriptional net-  
716 work. *Proceedings of the National Academy of Sciences* **100**:14796–14799. URL  
717 <http://www.pnas.org/cgi/doi/10.1073/pnas.2036429100>.
- 718 Kazanci, C., and Q. Ma, 2015. Chapter 3 System-wide measures in ecological net-  
719 work analysis. Pages 45–68 *in* Y.-S. Park, S. Lek, C. Baehr, and S. E. Jorgensen,  
720 editors. *Advanced Modelling Techniques Studying Global Changes in Environmen-*  
721 *tal Sciences*, volume 27. Elsevier.
- 722 Kéfi, S., E. L. Berlow, E. A. Wieters, S. A. Navarrete, O. L. Petchey, S. A. Wood,  
723 A. Boit, L. N. Joppa, K. D. Lafferty, R. J. Williams, N. D. Martinez, B. A.  
724 Menge, C. A. Blanchette, A. C. Iles, and U. Brose. 2012. More than a meal{...}  
725 integrating non-feeding interactions into food webs. *Ecology Letters* **15**:291–300.  
726 URL <http://doi.wiley.com/10.1111/j.1461-0248.2011.01732.x>.
- 727 Kones, J. K., K. Soetaert, D. van Oevelen, and J. O. Owino. 2009. Are network  
728 indices robust indicators of food web functioning? A Monte Carlo approach. *Eco-*  
729 *logical Modelling* **220**:370–382. URL [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/pii/S0304380008005024)  
730 [article/pii/S0304380008005024](http://www.sciencedirect.com/science/article/pii/S0304380008005024).
- 731 Krause, A. E., K. A. Frank, D. M. Mason, R. E. Ulanowicz, and W. W. Taylor.  
732 2003. Compartments revealed in food-web structure. *Nature* **426**:282–5. URL  
733 <http://dx.doi.org/10.1038/nature02115>.
- 734 Langfelder, P., and S. Horvath. 2008. WGCNA: an R package for weighted correlation  
735 network analysis. *BMC bioinformatics* **9**:559.
- 736 Lau, M. K., S. R. Borrett, D. E. Hines, and P. Singh, 2015. enaR: Tools for Ecological  
737 Network Analysis. URL <https://cran.r-project.org/web/packages/enaR/>.
- 738 Layton, A., B. Bras, and M. Weissburg. 2016. Ecological Principles and Metrics  
739 for Improving Material Cycling Structures in Manufacturing Networks. *Journal of*  
740 *Manufacturing Science and Engineering* **138**:101002.
- 741 Lédée, E. J. I., M. R. Heupel, A. J. Tobin, A. Mapleston, and C. A. Simpfendorfer.  
742 2016. Movement patterns of two carangid species in inshore habitats characterised  
743 using network analysis. *Marine Ecology Progress Series* **553**:219–232.
- 744 Legendre, P., L. Legendre, L. Legendre, and P. Legendre. 2012. *Numerical ecology*.  
745 Elsevier.

- 746 Lerner, B., and E. Boose, 2014. RDataTracker: Collecting Provenance in an In-  
 747 teractive Scripting Environment. Pages 1–4 *in* 6th USENIX Workshop on the  
 748 Theory and Practice of Provenance (TaPP 2014). USENIX Association, Cologne.  
 749 URL [https://www.usenix.org/conference/tapp2014/agenda/presentation/](https://www.usenix.org/conference/tapp2014/agenda/presentation/lerner)  
 750 [lerner](https://www.usenix.org/conference/tapp2014/agenda/presentation/lerner).
- 751 Lima, M. 2011. Visual Complexity: Mapping Patterns of Information. Princeton  
 752 Architectural Press.
- 753 Lindeman, R. L. 1942. The trophic-dynamic aspect of ecology. *Ecology* **23**:399–418.
- 754 Lowndes, J. S. S., B. D. Best, C. Scarborough, J. C. Afflerbach, M. R. Frazier, C. C.  
 755 O’Hara, N. Jiang, and B. S. Halpern. 2017. Our path to better science in less  
 756 time using open data science tools. *Nature Ecology & Evolution* **1**:0160. URL  
 757 <http://www.nature.com/articles/s41559-017-0160>.
- 758 Lubchenco, J., A. M. Olson, L. B. Brubaker, S. R. Carpenter, M. M. Hol-  
 759 land, S. P. Hubbell, S. A. Levin, J. A. MacMahon, P. A. Matson, J. M.  
 760 Melillo, H. A. Mooney, C. H. Peterson, and H. Ronald Pulliam. 1991. The  
 761 Sustainable Biosphere Initiative: An Ecological Research Agenda: A Report  
 762 from the Ecological Society of America. *Risser Source: Ecology* **72**:371–412.  
 763 URL <http://www.jstor.org/stable/2937183>[http://www.jstor.org/http://](http://www.jstor.org/http://www.jstor.org/action/showPublisher?publisherCode=esa)  
 764 [www.jstor.org/action/showPublisher?publisherCode=esa](http://www.jstor.org/http://www.jstor.org/action/showPublisher?publisherCode=esa).
- 765 Ludovisi, A., and U. M. Scharler. 2017. Towards a sounder interpretation of entropy-  
 766 based indicators in ecological network analysis. *Ecological Indicators* **72**:726–737.
- 767 MacArthur, R. 1955. Fluctuations of Animal Populations and a Measure of Com-  
 768 munity Stability. *Ecology* **36**:533. URL [http://www.readcube.com/articles/](http://www.readcube.com/articles/10.2307/1929601)  
 769 [10.2307/1929601](http://www.readcube.com/articles/10.2307/1929601).
- 770 Manly, B. F. J. 2007. Randomization, bootstrap and Monte Carlo methods in  
 771 biology. Chapman and Hall. URL [http://www.loc.gov/catdir/toc/fy0702/](http://www.loc.gov/catdir/toc/fy0702/2006047407.html)  
 772 [2006047407.html](http://www.loc.gov/catdir/toc/fy0702/2006047407.html).
- 773 Margalef, R. 1963. Certain unifying principles in ecology. *The American Naturalist*  
 774 **97**:357–374.
- 775 Martinez, N. D., 1992. Constant Connectance in Community Food Webs.



- 776 Maslov, S., and K. Sneppen. 2002. Specificity and stability in topology of protein  
777 networks. *Science* (New York, N.Y.) **296**:910–3. URL [http://www.ncbi.nlm.  
778 nih.gov/pubmed/11988575](http://www.ncbi.nlm.nih.gov/pubmed/11988575).
- 779 May, R. M. 1972. Will a Large Complex System be Stable? *Nature* **238**:413–414.  
780 URL <http://dx.doi.org/10.1038/238413a0>.
- 781 May, R. M. 2001. *Stability and Complexity in Model Ecosystems*. Princeton Uni-  
782 versity Press. URL [http://books.google.com/books?hl=en&lr=&id=  
783 BDA5-ipCLt4C&pgis=1](http://books.google.com/books?hl=en&lr=&id=BDA5-ipCLt4C&pgis=1).
- 784 May, R. M. 2006. Network structure and the biology of populations. *Trends in ecol-  
785 ogy {&} evolution* **21**:394–399. URL [http://www.ncbi.nlm.nih.gov/pubmed/  
786 16815438](http://www.ncbi.nlm.nih.gov/pubmed/16815438).
- 787 McNutt, M., K. Lehnert, B. Hanson, B. A. Nosek, A. M. Ellison, and J. L. King.  
788 2016. Liberating field science samples and data. *Science* **351**:1024–1026. URL  
789 <http://science.sciencemag.org/content/351/6277/1024.abstract>.
- 790 Milo, R., S. Shen-Orr, S. Itzkovitz, N. Kashtan, D. Chklovskii, U. Alon, S. H. Stro-  
791 gatz, D. Watts, S. H. Strogatz, A.-L. Barabási, R. Albert, M. Newman, H. Jeong,  
792 B. Tombor, R. Albert, Z. N. Oltvai, A. L. Barabasi, R. F. Cancho, C. Janssen,  
793 R. V. Sole, R. F. Cancho, R. V. Sole, L. Amaral, A. Scala, M. Barthélemy,  
794 H. Stanley, B. Huberman, L. Adamic, S. Shen-Orr, R. Milo, S. Mangan, U. Alon,  
795 N. Guelzim, S. Bottani, P. Bourguin, F. Kepes, M. Newman, S. H. Strogatz,  
796 D. Watts, S. Maslov, K. Sneppen, D. Thieffry, A. M. Huerta, E. Perez-Rueda,  
797 J. Collado-Vides, M. C. Costanzo, R. Williams, N. Martinez, S. Pimm, J. Law-  
798 ton, J. Cohen, J. White, E. Southgate, J. Thomson, S. Brenner, D. Callaway,  
799 J. Hopcroft, J. Kleinberg, M. Newman, and S. H. Strogatz. 2002. Network motifs:  
800 simple building blocks of complex networks. *Science* (New York, N.Y.) **298**:824–7.  
801 URL <http://www.ncbi.nlm.nih.gov/pubmed/12399590>.
- 802 Newman, M. 2010. *Networks an Introduction*. Oxford University  
803 Press. URL [http://www.oxfordscholarship.com/view/10.1093/acprof:oso/  
804 9780199206650.001.0001/acprof-9780199206650](http://www.oxfordscholarship.com/view/10.1093/acprof:oso/9780199206650.001.0001/acprof-9780199206650).
- 805 Newman, M. E. J., 2003. *The Structure and Function of Complex Networks*.
- 806 Newman, M. E. J. 2006. Modularity and community structure in networks. *Pro-  
807 ceedings of the National Academy of Sciences of the United States of America*  
808 **103**:8577–82. URL <http://www.pnas.org/content/103/23/8577.short>.

809 Newman, M. E. J., A. Clauset, C. Aicher, A. Z. Jacobs, A. Clauset, S. Fortu-  
 810 nato, P. Holme, M. Huss, H. Jeong, R. Guimerà, L. A. N. Amaral, D. Hric,  
 811 R. K. Darst, S. Fortunato, M. Barthélemy, A. Z. Jacobs, A. Clauset, K. Zuev,  
 812 G. B. M. Boguñá, D. Krioukov, B. H. Good, Y.-A. d. Montjoye, A. Clauset,  
 813 C. Bothorel, J. D. Cruz, M. Magnani, B. Micenková, N. Binkiewicz, J. T. Vogel-  
 814 stein, K. Rohe, E. Galbrun, A. Gionis, N. Tatti, T. J. Hansen, M. W. Mahoney,  
 815 Y. Zhang, E. Levina, J. Zhu, P. Expert, T. S. Evans, V. D. Blondel, R. Lambiotte,  
 816 L. Peel, P. Zhang, C. Moore, L. Zdeborová, X. Maa, L. Gaoa, X. Yongb, L. Fua,  
 817 Z.-Y. Zhang, P. W. Holland, K. B. Laskey, S. Leinhardt, B. Karrer, M. E. J. New-  
 818 man, A. Decelle, F. Krzakala, C. Moore, L. Zdeborová, E. Mossel, J. Neeman,  
 819 A. Sly, J. Moody, L. Danon, J. Duch, A. Diaz-Guilera, A. Arenas, A. F. Mc-  
 820 Daid, D. Greene, N. Hurley, U. Brose, G. Woodward, A. L. Traud, P. J. Mucha,  
 821 M. A. Porter, P. C. Bull, D. B. Larremore, A. Clauset, C. Z. Buckee, D. B.  
 822 Larremore, A. Clauset, A. Z. Jacobs, G. M. Warimwe, P. C. Bull, A. Clauset,  
 823 C. Moore, M. E. J. Newman, R. Milo, S. P. Borgatti, M. G. Everett, B. Ball,  
 824 M. E. J. Newman, P. D. Hoff, A. E. Raftery, and M. S. Handcock. 2016. Struc-  
 825 ture and inference in annotated networks. *Nature Communications* **7**:11863. URL  
 826 <http://www.nature.com/doifinder/10.1038/ncomms11863>.

827 Newman, M. E. J., S. H. Strogatz, and D. J. Watts. 2001. Random graphs with  
 828 arbitrary degree distributions and their applications. *Physical Review E* **64**:026118.  
 829 URL <http://arxiv.org/abs/cond-mat/0007235/>.

830 Noble, W. S. 2009. A quick guide to organizing computational biology projects.  
 831 *PLoS computational biology* **5**:e1000424. URL [http://dx.plos.org/10.1371/](http://dx.plos.org/10.1371/journal.pcbi.1000424)  
 832 [journal.pcbi.1000424](http://dx.plos.org/10.1371/journal.pcbi.1000424).

833 Odum, H. T. 1957. Trophic structure and productivity of Silver Springs, Florida.  
 834 *Ecol. Mono.* **27**:55–112.

835 Odum, H. T., and R. C. Pinkerton. 1955. Time’s speed regulator: the optimum  
 836 efficiency for maximum power output in physical and biological systems. *American*  
 837 *Scientist* **43**:331–343.

838 Paine, R. T. 1966. Food Web Complexity and Species Diversity. *The American*  
 839 *Naturalist* **100**:65.

840 Parker, T. H., S. Nakagawa, and J. Gurevitch. 2016. Promoting transparency in  
 841 evolutionary biology and ecology. *Ecology Letters* **19**:726–728. URL [http://](http://doi.wiley.com/10.1111/ele.12610)  
 842 [doi.wiley.com/10.1111/ele.12610](http://doi.wiley.com/10.1111/ele.12610).

- 843 Patten, B. C. 1978. Systems approach to the concept of environment. *Ohio Journal*  
844 *of Science* **78**:206–222. URL <https://kb.osu.edu/dspace/handle/1811/22549>.
- 845 Patten, B. C., and G. T. Auble. 1981. System theory of the ecological niche. *American*  
846 *Naturalist* **117**:893–922. URL <GotoISI>://BIOABS:BACD198172072401.
- 847 Patten, B. C., and M. Witkamp. 1967. Systems analysis of  $^{134}\text{cesium}$  kinetics in  
848 terrestrial microcosms. *Ecology* **48**:813–824.
- 849 Petanidou, T., A. S. Kallimanis, J. Tzanopoulos, S. P. Sgardelis, and J. D. Pantis.  
850 2008. Long-term observation of a pollination network: Fluctuation in species and  
851 interactions, relative invariance of network structure and implications for estimates  
852 of specialization. *Ecology Letters* **11**:564–575.
- 853 Pimm, S. L. 1982. Food webs. Chapman and Hall, London; New York.
- 854 Poisot, T., B. Baiser, J. A. Dunne, S. Kéfi, F. Massol, N. Mouquet, T. N. Ro-  
855 manuk, D. B. Stouffer, S. A. Wood, and D. Gravel. 2015. mangal mak-  
856 ing ecological network analysis simple. *Ecography* pages n/a–n/a. URL <http://onlinelibrary.wiley.com/doi/10.1111/ecog.00976/abstract>.  
857
- 858 Poisot, T., D. Gravel, S. Leroux, S. A. Wood, M.-J. Fortin, B. Baiser, A. R. Cirtwill,  
859 M. B. Araújo, and D. B. Stouffer. 2016*a*. Synthetic datasets and community  
860 tools for the rapid testing of ecological hypotheses. *Ecography* **39**:402–408. URL  
861 <http://doi.wiley.com/10.1111/ecog.01941>.
- 862 Poisot, T., D. B. Stouffer, and S. Kéfi. 2016*b*. Describe, understand and predict:  
863 why do we need networks in ecology? *Functional Ecology* **30**:1878–1882. URL  
864 <http://doi.wiley.com/10.1111/1365-2435.12799>.
- 865 Post, D. M., M. L. Pace, and N. G. Hairston. 2000. Ecosystem size determines  
866 food-chain length in lakes. *Nature* **405**:1047–1049.
- 867 Proulx, S. R., D. E. L. Promislow, and P. C. Phillips. 2005. Network thinking in  
868 ecology and evolution. *Trends in ecology & evolution* **20**:345–53. URL <http://www.ncbi.nlm.nih.gov/pubmed/16701391>.  
869
- 870 QUINTESENCE Consortium, M. E., S. Carpenter, e. al., G. Daily, e. al.,  
871 C. Raudsepp-Hearne, e. al., I. P. o. B. Services, Ecosystem, J. Silvertown,  
872 M. Schröter, e. al., S. Naeem, e. al., N. Biggs, e. al., A. Mashaghi, e. al.,

873 T. Ideker, R. Sharan, J. Gardy, e. al., T. Valente, D. Acemoglu, e. al., M. Jack-  
874 son, M. Janssen, e. al., G. Woodward, e. al., C. Mulder, J. Elser, R. Thomp-  
875 son, e. al., J. Montoya, e. al., O. Petchey, e. al., J. Reiss, e. al., A. Barabasi,  
876 R. Albert, F. Harary, R. Haines-Young, S. Macfadyen, e. al., K. Rathwell,  
877 G. Peterson, R. Costanza, I. Kubiszewski, C. Mulder, e. al., H. Tallis, e. al.,  
878 A. Tavoni, S. Levin, A. Ma, R. Mondragón, M. Pascual, J. Dunne, F. Jordán,  
879 e. al., H. Ernstson, e. al., U. Narloch, e. al., M. Hagen, e. al., R. Stewart,  
880 e. al., A. Shmida, M. Wilson, M. Palmer, e. al., G. Mace, e. al., R. d. Groot,  
881 e. al., H. Zimmermann, R. Albert, e. al., M. Pocock, e. al., K. Chan, e. al.,  
882 L. Dicks, e. al., G. McInerny, e. al., M. Pocock, e. al., A. Vespignani, D. Bo-  
883 han, e. al., P. Anderson, J. Cohen, e. al., J. Montoya, e. al., S. Pimm, C. Holling,  
884 A. Trichard, e. al., D. Bohan, and e. al. 2016. Networking Our Way to Better  
885 Ecosystem Service Provision. *Trends in ecology & evolution* **31**:105–15. URL  
886 <http://www.ncbi.nlm.nih.gov/pubmed/26777789>.

887 R Core Team, 2017. R: A Language and Environment for Statistical Computing.  
888 R Foundation for Statistical Computing, Vienna, Austria. URL [https://www.](https://www.R-project.org/)  
889 [R-project.org/](https://www.R-project.org/).

890 Samuelson, P. A. 1948. *Economics: An Introductory Analysis*. McGraw–Hill Book  
891 Co., New York,.

892 Shen, H.-W., and A.-L. Barabasi. 2014. Collective credit allocation in science.  
893 *Proceedings of the National Academy of Sciences* **111**:12325–12330. URL [http:](http://www.pnas.org/content/111/34/12325.abstract)  
894 [//www.pnas.org/content/111/34/12325.abstract](http://www.pnas.org/content/111/34/12325.abstract).

895 Sih, A., S. F. Hanser, and K. a. McHugh. 2009. Social network theory: New insights  
896 and issues for behavioral ecologists. *Behavioral Ecology and Sociobiology* **63**:975–  
897 988.

898 Simberloff, D., and B. V. Holle. 1999. Positive interactions of nonindigenous species:  
899 invasional meltdown? *Biological Invasions* pages 21–32.

900 Solé, R. V., and J. M. Montoya. 2001. Complexity and fragility in ecological networks.  
901 *Proceedings of the Royal Society B* **268**:2039–2045.

902 Stouffer, D. B., and J. Bascompte. 2010. Understanding food-web persistence from  
903 local to global scales. *Ecology Letters* **13**:154–161. URL [http://doi.wiley.com/](http://doi.wiley.com/10.1111/j.1461-0248.2009.01407.x)  
904 [10.1111/j.1461-0248.2009.01407.x](http://doi.wiley.com/10.1111/j.1461-0248.2009.01407.x).

- 905 Strogatz, S. H. 2001. Exploring complex networks. *Nature* **410**:268–76. URL  
906 <http://dx.doi.org/10.1038/35065725>.
- 907 Stumpf, M. P. H., W. P. Kelly, T. Thorne, and C. Wiuf. 2007. Evolution at the system  
908 level: the natural history of protein interaction networks. *Trends in ecology &  
909 evolution* **22**:366–373. URL <http://www.ncbi.nlm.nih.gov/pubmed/17475365>.
- 910 Sutherland, W. J., R. P. Freckleton, H. C. J. Godfray, S. R. Beissinger, T. Ben-  
911 ton, D. D. Cameron, Y. Carmel, D. A. Coomes, T. Coulson, M. C. Emmer-  
912 son, R. S. Hails, G. C. Hays, D. J. Hodgson, M. J. Hutchings, D. Johnson,  
913 J. P. G. Jones, M. J. Keeling, H. Kokko, W. E. Kunin, X. Lambin, O. T. Lewis,  
914 Y. Malhi, N. Mieszkowska, E. J. Milner-Gulland, K. Norris, A. B. Phillimore,  
915 D. W. Purves, J. M. Reid, D. C. Reuman, K. Thompson, J. M. J. Travis,  
916 L. A. Turnbull, D. A. Wardle, and T. Wiegand. 2013. Identification of 100  
917 fundamental ecological questions. *Journal of Ecology* **101**:58–67. URL <http://doi.wiley.com/10.1111/1365-2745.12025>.
- 919 Tansley, A. G. 1935. The Use and Abuse of Vegetational Concepts and Terms.  
920 *Ecology* **16**:284–307. URL [http://links.jstor.org/sici?sici=0012-9658\(193507\)29:1<28193507:3A333C2843ATUAA0V3E2.0.CO;3B2-P](http://links.jstor.org/sici?sici=0012-9658(193507)29:1<28193507:3A333C2843ATUAA0V3E2.0.CO;3B2-P).
- 922 Ulanowicz, R. E., 1986. Introduction. Pages 1–8 *in* Growth and Develop-  
923 ment. Springer New York, New York, NY. URL [http://link.springer.com/10.1007/978-1-4612-4916-0\\_1](http://link.springer.com/10.1007/978-1-4612-4916-0_1)  
924 [http://link.springer.com/10.1007/](http://link.springer.com/10.1007/978-1-4612-4916-0_1)  
925 [978-1-4612-4916-0\\_1](http://link.springer.com/10.1007/978-1-4612-4916-0_1).
- 926 Ulanowicz, R. E., R. D. Holt, and M. Barfield. 2014. Limits on ecosystem trophic  
927 complexity: insights from ecological network analysis. *Ecology letters* **17**:127–36.  
928 URL <http://www.ncbi.nlm.nih.gov/pubmed/24382355>.
- 929 Ulrich, W., and N. J. Gotelli. 2007. Null model analysis of species nestedness  
930 patterns. *Ecology* **88**:1824–1831.
- 931 Ulrich, W., and N. J. Gotelli. 2010. Null model analysis of species associations  
932 using abundance data. *Ecology* **91**:3384–97. URL <http://www.ncbi.nlm.nih.gov/pubmed/21141199>.
- 934 Urban, D., and T. Keitt. 2001. Landscape connectivity: A graph-theoretic perspec-  
935 tive. *Ecology* **82**:1205–1218.
- 936 Valente, T. W., K. Coronges, C. Lakon, and E. Costenbader. 2008. How Correlated  
937 Are Network Centrality Measures? *Connections (Toronto, Ont.)* **28**:16–26.

- 938 Vermaat, J. E., J. a. Dunne, and A. J. Gilbert. 2009. Major dimensions in food-web  
939 structure properties. *Ecology* **90**:278–282. URL [http://www.ncbi.nlm.nih.gov/  
940 pubmed/19294932](http://www.ncbi.nlm.nih.gov/pubmed/19294932).
- 941 Visser, M. D., S. M. McMahon, C. Merow, P. M. Dixon, S. Record, E. Jonge-  
942 jans, W. Michener, M. Jones, S. Petrovskii, N. Petrovskaya, A. Ellison, B. Den-  
943 nis, G. Hager, G. Wellein, P. Zuidema, E. Jongejans, P. Chien, H. During,  
944 F. Schieving, B. V. Putten, M. Visser, H. Muller-Landau, P. Jansen, L. Comita,  
945 H. Muller-Landau, S. Aguilar, S. Hubbell, C. Merow, N. LaFleur, J. Silander,  
946 A. Wilson, M. Rubega, M. Visser, E. Jongejans, M. V. Breugel, P. Zuidema,  
947 Y.-Y. Chen, A. Kassim, J. Chambers, B. Kernighan, P. Plauger, G. Amdahl,  
948 A. Porter, R. Selby, R. Bryant, D. OHallaron, M. Schmidberger, M. Morgan,  
949 D. Eddelbuettel, H. Yu, L. Tierney, U. Mansmann, A. Grama, G. Karypis,  
950 V. Kumar, A. Gupta, P. LEcuyer, D. Eddelbuettel, R. François, D. Eddelbuet-  
951 tel, C. Sanderson, H. Robbins, S. Monro, A. Mantoglou, J. Wilson, A. Finley,  
952 Z. Merali, A. Guisan, A. Lehmann, S. Ferrier, M. Austin, J. Overton, B. Brook,  
953 J. OGrady, A. Chapman, M. Burgman, H. Akçakaya, R. Frankham, F. Isbell,  
954 V. Calcagno, A. Hector, J. Connolly, W. Harpole, P. Reich, E. V. Moran, J. Clark,  
955 G. Bohrer, G. Katul, R. Walko, and R. Avissar. 2015. Speeding Up Eco-  
956 logical and Evolutionary Computations in R; Essentials of High Performance  
957 Computing for Biologists. *PLOS Computational Biology* **11**:e1004140. URL  
958 <http://dx.plos.org/10.1371/journal.pcbi.1004140>.
- 959 Wang, S., and B. Chen. 2016. Energy-water nexus of urban agglomeration based on  
960 multiregional input-output tables and ecological network analysis: A case study  
961 of the Beijing-Tianjin-Hebei region. *Applied Energy* **178**:773–783.
- 962 Wasserman, S., and K. Faust. 1994. *Advances in Social Network Analysis: Research*  
963 *in the Social and Behavioral Sciences*. SAGE Publications. URL [http://books.  
964 google.com/books?hl=en&lr=&id=C6juDKDmvCcC&pgis=1](http://books.google.com/books?hl=en&lr=&id=C6juDKDmvCcC&pgis=1).
- 965 Williams, R. J., E. L. Berlow, J. a. Dunne, A.-L. Barabási, and N. D. Mar-  
966 tinez. 2002. Two degrees of separation in complex food webs. *Proceed-*  
967 *ings of the National Academy of Sciences of the United States of America*  
968 **99**:12913–12916. URL [http://www.pubmedcentral.nih.gov/articlerender.  
969 fcgi?artid=130559&tool=pmcentrez&rendertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=130559&tool=pmcentrez&rendertype=abstract).
- 970 Williams, R. J., and N. D. Martinez. 2000. Simple rules yield complex food  
971 webs. *Nature* **404**:180–183. URL [http://www.nature.com/doifinder/10.1038/  
972 35004572](http://www.nature.com/doifinder/10.1038/35004572).

- 973 Wilson, E. O., 1999. Consilience: The unity of knowledge.
- 974 Woodward, G. U. Y., J. P. Benstead, O. S. Beveridge, J. Blanchard, T. Brey, L. E.  
 975 E. E. Brown, W. F. Cross, N. Friberg, C. Ings, U. T. E. Jacob, S. Jennings, M. E.  
 976 Ledger, A. M. Milner, J. M. Montoya, E. O. Gorman, J. M. Olesen, O. L. Petchey,  
 977 E. Pichler, D. C. Reuman, M. S. A. Thompson, F. J. F. V. a. N. Veen, G. Yvon-  
 978 Durocher, T. C. Ings, U. T. E. Jacob, S. Jennings, M. E. Ledger, A. M. Milner,  
 979 J. M. Montoya, E. O 'gorman, J. M. Olesen, O. L. Petchey, D. E. Pichler, D. C.  
 980 Reuman, M. S. A. Thompson, F. J. F. Van Veen, and G. Yvon-Durocher. 2010.  
 981 Ecological Networks in a Changing Climate. *Advances In Ecological Research* 42.  
 982 Ecological Networks. **42**:72–120.
- 983 Xia, L., B. D. Fath, U. M. Scharler, and Y. Zhang. 2016. Spatial variation in the  
 984 ecological relationships among the components of {Beijing}'s carbon metabolic  
 985 system. *Science of the Total Environment* **544**:103–113.
- 986 Yang, J., and B. Chen. 2016. Energy–water nexus of wind power generation systems.  
 987 *Applied Energy* **169**:1–13.
- 988 Yodzis, P., and S. Innes. 1992. Body Size and Consumer-Resource Dynamics. *The*  
 989 *American Naturalist* **139**:1151–1175. URL <http://www.journals.uchicago.edu/doi/10.1086/285380>.
- 991 Zhao, L., H. Zhang, E. J. O’Gorman, W. Tian, A. Ma, J. C. Moore, S. R. Borrett,  
 992 and G. Woodward. 2016. Weighting and indirect effects identify keystone species  
 993 in food webs. *Ecology Letters* **19**:1032–1040.
- 994 Zinkgraf, M., L. Liu, A. Groover, and V. Filkov. 2017. Identifying gene coex-  
 995 pression networks underlying the dynamic regulation of wood-forming tissues in  
 996 *Populus* under diverse environmental conditions. *New Phytologist* URL  
 997 <http://doi.wiley.com/10.1111/nph.14492>.

## 998 Acknowledgments

999 This work was supported by the US National Science Foundation under grant SSI-  
 1000 1450277 End-to-End Provenance

## <sup>1001</sup> **Author contributions statement**

<sup>1002</sup> All authors contributed to the conception, writing and review of the manuscript.



## Boxes

*Box 1. Benchmarking Ecological Models* The most basic test is to feed the algorithm a set of "random" matrices to make sure that the frequency of statistically significant results is no greater than 5%. Otherwise, the algorithm is vulnerable to a Type I statistical error (incorrectly rejecting a true null hypothesis). However, specifying a matrix produced by random sampling errors is not so easy. By definition, if a null model algorithm is used to generate the random matrices, then no more than 5% of them should be statistically significant (unless there were programming errors). For binary matrices, two log-normal distributions can be used to generate realistic heterogeneity in row and column totals, while still maintaining additive effects for cell occurrence probabilities (Ulrich and Gotelli, 2010). "Structured" matrices are needed to test for Type II errors (incorrectly accepting a false null hypothesis), and these require a careful consideration of exactly what sort of pattern or mechanism the test is designed to reveal. One approach is to begin with a perfectly structured matrix, such as one derived from a mechanistic model for generating network structure, contaminate it with increasing amounts of stochastic noise, and test for the statistical pattern at each step (Gotelli, 2000). A plot of the  $P$  value versus the added noise should reveal an increasing curve, and will indicate the signal-to-noise ratio below which the test cannot distinguish the pattern from randomness. Alternatively, one can begin with a purely random matrix but embed in it a non-random substructure, such as a matrix clique or a node with extreme centrality. The size, density, and other attributes of this matrix can be manipulated to see whether the test can still detect the presence of the embedded structure (Gotelli et al., 2010). Because all null model tests (and all frequentist statistics) are affected by sample size and data structure, these benchmark tests can be tailored to the attributes of the empirical data structures for better focus and improved inference.

Even simple randomization algorithms may require further filters to ensure that random matrices retain a number of desirable network properties. For example, Dunne et al. (2002) created random food-web matrices with constant species richness and connectance, but they discarded webs with unconnected nodes and subwebs because these topologies were not observed in the empirical webs. A "stub reconstruction" algorithm builds a topology that is constrained to the observed number of edges per node (Newman et al., 2001). Each node is assigned the correct number of edges, and then nodes are successively and randomly paired to create a growing network. However, this algorithm also generates multiple edges between the same two nodes, which must be discarded or otherwise accounted for. Maslov and Sneppen (2002) use a "local re-wiring algorithm" that preserves the number of connections

1040 for every node by swapping edges randomly between different pairs of nodes. This  
1041 algorithm is closely analogous to the swap algorithm used in species co-occurrence  
1042 analyses that preserves the row and column totals of the original matrix (Connor  
1043 and Simberloff, 1979). The more constraints that are added to the algorithm, the  
1044 less likely it is that simple sampling processes can account for patterns in the data.  
1045 However, some constraints, such as connectivity or matrix density, may inadvertently  
1046 “smuggle in” the very processes they are designed to detect. This can lead to the  
1047 so-called “Narcissus” effect (Colwell and Winkler, 1984). Finding the correct balance  
1048 between realistic constraints and statistical power is not easy (Gotelli et al., 2012),  
1049 and there are many potential algorithms that reasonably could be used, even for  
1050 simple binary matrices (Gotelli, 2000).

# Tables

Sub-discipline	Level	Metric	Concept	Reference
General	W	Density	The proportion of possible edges that are actually associated with nodes; called Connectance in Food Web ecology.	
General	N	Centrality	Multiple ways to characterize the relative importance of nodes.	Wasserman and Faust (1994)
General	N	Degree	Number of edges connected to a given node, which is a type of local centrality.	
General	N	Eigenvector Centrality	Global centrality metric based on number of walks that travel through a node	Bonacich (1987)
General	W	Centrality Distribution	Shape of the frequency distribution of edges among nodes.	Barabási and Albert (1999); Dunne et al. (2002)
General	W	Centralization	The concentration (versus evenness) of centrality among the nodes.	Freeman (1979)
General	W	Graph diameter	The longest path between any two nodes in a graph.	Barabási et al. (2000); Urban and Keitt (2001)
General	W	Modularity	Degree to which edges are distributed within rather than between distinct sets of nodes.	Newman (2010)
General	G	Motifs	Small sets of nodes with similar distributions of edges.	Milo et al. (2002)
General	W	Link density	Average number of edges per node.	Martinez (1992)
Community	N	Temperature	Measures the nestedness of a bipartite network.	Ulrich and Gotelli (2007)
Community	W	Co-occurrence	Degree of overlapping spatial or temporal distributions of species relative to a null model.	Gotelli (2000)
Community	N	Indicator Species	Degree to which the abundance of a taxonomic group responds to an environmental gradient.	
Community	W	Nestedness	The degree to which interactions can be arranged into subsets of the larger community	
Community	W	Evenness	Deviation of the distribution of observed abundances relative to an even distribution among taxonomic groups in a community	
Community	W	Diversity	Distribution of abundances among taxonomic groups in an observed community	
Community	W	Richness	The number of taxonomic groups in a community	
Community	W	Stability	The change in the abundances of taxonomic groups across a set of observations	
Food-Web	N	Removal Importance	The degree to which removal of a compartment or species produces subsequent removals in the ecosystem.	Borrvall et al. (2000); Dunne et al. (2002); Eklöf and Ebenman (2006); Solé and Montoya (2001)
General	N	Connectance	Proportion of realized out of possible edges	Pimm (1982); Vernaas et al. (2009)
Food-Web	G	Food-chain length	The number of feeding relationships among a set of compartments in a food-web.	Post et al. (2000); Ulanowicz et al. (2014)
Ecosystem	W	Finn cycling index	Degree to which matter or energy passes through the same set of compartments.	Finn (1980)
Ecosystem	G	Environ	The sub-network of the probability of movement of energy or matter among compartments generated by a single unit of input (output) into a selected node.	Patten (1978); Patten and Auble (1981)
Ecosystem	N	Throughflow	Amount of energy or matter passing into or out of a node	Finn (1976)
Ecosystem	N	Throughflow Centrality	The proportion of energy or matter that passes through a given compartment in an ecosystem.	Borrett (2013)
General	G	Chain Length	Number of edges between two nodes in a group	
Food-Web	G	Average Path Length	The average number of times a unit of matter or energy travels from one compartment to another before exiting the ecosystem	Finn (1976)
Ecosystem	W	Pathway Proliferation	Rate of increase in the number of edges between nodes with increasing path length.	Borrett et al. (2007)
Ecosystem	W	Ascendency	Measures the average similarity in matter or energy flows among compartments in an ecosystem.	Ulanowicz (1986)
Food-Web	N	Trophic Level	Ordinal classification of a compartment or taxonomic group based on the relative position in the ecosystem.	Allesina and Pascual (2009); Fath (2004); Williams et al. (2002)

Table 1: Ecological network metric summary and classification. Level indicates the hierarchy of the metric (W = Whole network, G = Group or sub-network, N = Node). The Sub-disciplines include 'General' network theory, 'Community' ecology, 'Food-web' and 'Ecosystem' ecology. Also available at <https://figshare.com/s/1bf1a7e0a6ee3ac97a4b>

# Figures




Figure 1: Although systems thinking has been a part of ecology since at least the work of Darwin, network ecology has grown rapidly since the turn of the last century but has been developing in isolated sub-fields. (A) Plot showing the increase in “network ecology” keywords in the literature from 1991 to current (updated search based on Borrett et al., 2014). (B) Contour plot of common topics in network ecology with peaks indicating clusters of related topics. The regions are labeled with the most common terms found in the clusters. From Borrett et al. (2014), reproduced with permission.




Figure 2: Hypothetical unweighted, directed network showing examples of the four classes of network metrics. *Node Level:* the purple node exhibits low centrality while the orange node exhibits high centrality. *Group or Sub-Network Level:* the blue nodes connected with dashed edges shows a module. *Global or Whole Network Level:* using the edges of all nodes we can measure the connectance of the entire network ( $c = \text{edges}/\text{nodes}^2 = 0.12$ ).